



Review Article



Exploring Thermostable Lipases: Molecular Innovations and Expanding Industrial Horizons

Tehmina Bashir¹, Alamgir², Syeda Amna Batool³, Tahir Hussain⁴, Washdev⁵ and Iram Rafique⁶¹Department of Botany, Government College University, Lahore, Pakistan²Faculty of Allied Health Sciences, Sohail University, Karachi, Pakistan³Department of Botany, University of Narowal, Narowal, Pakistan⁴Faculty of Allied Health Sciences, Hamdard Institute of Management Sciences, Karachi, Pakistan⁵Chughtai Laboratory, Karachi, Pakistan⁶Faculty of Allied Health Sciences, Karachi Institute of Nursing Allied Health Sciences, Karachi, Pakistan

ARTICLE INFO

Keywords:

Thermo-Stable Lipases, Commercial Uses, Characterization, Enzyme Engineering, Biocatalysts

How to Cite:

Bashir, T., Alamgir, ., Batool, S. A., Hussain, T., Washdev, ., & Rafique, I. (2025). Exploring Thermostable Lipases: Molecular Innovations and Expanding Industrial Horizons: Exploring Thermostable Lipases: Molecular Innovations . Futuristic Biotechnology, 5(2), 28-35. <https://doi.org/10.54393/fbt.v5i2.183>

*Corresponding Author:

Tehmina Bashir

Department of Botany, Government College University, Lahore, Pakistan

tehmabashir25@gmail.com

Received Date: 23rd April, 2025Revised Date: 20th June, 2025Acceptance Date: 25th June, 2025Published Date: 30th June, 2025

ABSTRACT

Thermostable lipases are crucial biocatalysts valued for their stability and functionality in both aqueous and non-aqueous environments, enabling efficient catalysis at elevated temperatures. These enzymes, derived from both wild-type and genetically engineered strains, exhibit unique properties that make them indispensable across diverse industrial sectors. Despite their potential, challenges remain in optimizing their production, purification, and characterization to meet specific application requirements ranging from highly purified pharmaceutical formulations to less refined industrial uses. This review consolidates recent advances in the isolation, engineering, and detailed characterization of thermostable lipases, highlighting their substrate specificity, catalytic efficiency, enantioselectivity, and tolerance to harsh conditions. Emphasis is placed on emerging molecular innovations and metagenomic approaches for discovering novel enzymes with enhanced industrial applicability. By bridging fundamental insights with practical applications, this overview aims to guide future research and development efforts in harnessing thermostable lipases for expanding biotechnological horizons.

INTRODUCTION

Thermostable lipases are extensively used in various biocatalytic reactions due to their excellent performance at elevated temperatures. Operating at high temperatures offers several advantages, including increased product formation rates with minimal diffusional limitations, improved conversion efficiency, enhanced solubility of hydrophobic substrates, greater molecular mobility, and a reduced risk of microbial contamination [1]. Lipases (EC 3.1.1.3, triacylglycerol acylhydrolases) rank among the most widely employed enzymes in organic synthesis and

biotechnological processes. Although lipases from extremophiles exhibit activity across diverse pH ranges, their stability often declines above 70°C. Typically, microbial lipases show optimal activity between 30°C and 65°C. Extremophilic organisms inhabiting various environments have been identified as sources of a broad spectrum of thermostable lipases, whose stability under limited water conditions enhances their industrial applicability [2, 3]. Many thermostable lipases have been characterized from moderate extremophiles, particularly



from the genus *Bacillus*, such as *Bacillus* sp. J-33 [4], *Lactobacillus acidophilus* [5], *Bacillus thermoleovorans* ID1 [6], and *Pseudomonas aeruginosa* species [7]. Advances in genomics and protein engineering continue to deepen our understanding of thermostable lipases. These enzymes maintain exceptional stability in both aqueous and non-aqueous environments at elevated temperatures, facilitating higher reaction rates, reducing medium viscosity, and minimizing microbial contamination [1, 2]. Recent studies highlight structural features that contribute to their enhanced properties, including robust hydrogen bonding networks, increased hydrophobic interactions, and optimized surface loops—traits commonly observed in enzymes from extremophiles. For example, semi-rational design strategies have been used to improve thermostability, as demonstrated in novel esterase structures [8]. The exploration for novel thermostable lipases increasingly employs metagenomic approaches, enabling the discovery of enzymes from unculturable microorganisms in diverse extreme environments, such as medicinal wastewater [9, 10]. This has expanded opportunities to isolate wild strains with unique features and to engineer recombinant strains tailored to biotechnological needs. The intrinsic heat resistance of proteins in extremophiles arises from multiple molecular interactions that stabilize their structure, including strengthened ionic networks, enhanced hydrophobic contacts, increased hydrogen bonding, the presence of disulfide bridges, shortened surface loops, and the capping of N- and C-termini. Moreover, thermophilic enzymes exhibit a densely packed and rigid framework that supports stability at high temperatures but often compromises catalytic activity at lower temperatures. This reduced activity at moderate temperatures limits their application in processes where fine chemicals remain stable only under such conditions. Additionally, these biocatalysts can be sensitive to adverse factors like organic solvents, ionic strength variations, extreme pH, high pressure, and temperature fluctuations, which may challenge their economic viability in industrial processes [11]. Thermal Stability (T_{max}), Optimum pH, and half-life ($t_{1/2}$) of thermostable lipases from various naturally sourced microorganisms, illustrating their diverse stability profiles at different temperatures, are shown in Table 1.

Table 1: Thermal Stability, Optimum pH, and Half-Life of Thermostable Lipases From Various Naturally Sourced Microorganisms

Source	T max°C	pH optimum	Stability	References
<i>Bacillaceae</i> sp. RSJ1	50°C	8-9	$t_{1/2}$ = 150, 90, 55, 7 and 45 min at 60°C, 65°C, 71°C and 76°C, respectively	[11]

<i>Pseudomonas</i> spp.	90°C	11	$t_{1/2}$ = 13 hr at 90°C	[12]
<i>Bacillus thermoleovorans</i> ID-1	70-75°C	7.5	$t_{1/2}$ = 30 min at 70°C and 1 hr at 60°C	[13]
<i>Bacillus</i> strain A30-1	60°C	6-9	$t_{1/2}$ = 8 hr at 75°C	[14]
<i>Aneurinibacillus thermoaerophilus</i> strain HZ	65°C	7	$t_{1/2}$ = 3 hr 10 min and 1 hr 20 min at 65°C and 70°C, respectively	[15]
<i>Burkholderia multivorans</i> V2	45-50°C	8	$t_{1/2}$ = ten and 50 minutes at fifty to sixty degrees Celsius, respectively	[16]
<i>Burkholderia</i> spp.	60°C	8.5	$t_{1/2}$ = 2 and 0.5 hr at 50°C and 60°C, respectively	[17]
<i>Bacillus subtilis</i> NS 8	60°C	7	$t_{1/2}$ = 4 hr 33 min at 60°C, 52 min at 71°C and ~42 min at 80°C	[18]
<i>Bacillus coagulans</i> MTCC-6375	45°C	8.5	$t_{1/2}$ = 4 hr 33 min at 60°C, 51 min at 70°C and ~42 min at 80°C	[19]
<i>Geobacillus thermodenitrificans</i> IBRL-nra	65°C	7	$t_{1/2}$ = 20 min at 55°C. 65°C 7 $t_{1/2}$ = 8 hr at 60°C, 16 hr at 65°C - 75°C	[20]

Optimum pH and temperature conditions for catalytic activity of thermostable lipases from various microorganisms are shown in Table 2.

Table 2: Optimum pH and Temperature Conditions for Catalytic Activity of Thermostable Lipases from Various Microorganisms

Microorganism	Enzyme Optimum pH	Enzyme Optimum Temperature (°C)	References
<i>Bacillaceae</i> sp. RSJ-1	8-9	50°C	[11]
<i>Bacillaceae acidocaldarius</i>	-	68 to 70°C	[21]
<i>Bacillaceae</i> strain J33	8	60°C	[22]
<i>Bacillus thermocatenulatus</i> BTL2	9	60-70°C	[23]
<i>Geobacillus Stearothermophilus</i>	-	65°C	[24]
<i>Geobacillus Stearothermophilus</i>	8	70°C	[25]
<i>Pseudomonas</i> spp.	15	90°C	[26]
<i>Pseudomonas</i> sp.	9.7	60°C	[27]
<i>Pyrobaculum calidifontis</i>	-	95°C	[28]
<i>Pyrococcus furiosus</i>	-	99°C	[29]

<i>Pyrococcus horikoshii</i>	7	95°C	[30]
<i>Pyrococcus horikoshii</i>	5.9	90°C	[31]
<i>Staphylococcus aureus</i>	5-12	55°C	[32]
<i>Rhizopus chinensis</i>	9	30-50°C	[33]
<i>Thermoanaerobacter thermohydrosulfuricus</i>	7-8	75-90°C	[34]
<i>Staphylococcus xylosus</i>	5-12	65-70°C	[35]
<i>Bacillus Subtilis</i>	10	55 to 79°C	[36]
<i>Bacillus pumilus</i> Rk31	7	40-70°C	[11]

Industrial Use of Thermostable Lipases

The application of substance catalysts doesn't just include many drawbacks linked to it; however as well provides an increase to several unwanted byproducts together with poisonous effluents, while at the same time, substrate uniqueness, bio-degradability, and lots of these kinds of positive aspects related to biocatalysts provide them with an advantage over the usage of catalysts in the chemical industrial sectors [37]. Thermostable enzymes can take on higher temperatures, therefore endowing a long half-life to the bio-catalyst. These types of enzymes may also put up with elevated levels of substrates as well as be protected from chemical denaturants. Their potential to perform different reactions at a high reaction rate as a result of a rise in substrate diffusion coefficient, as well as lower viscosity at elevated temperature ranges, make them a well-liked option over the sources of mesophiles [11]. The bio-technological possibilities related to lipases cause them to become a good biocatalyst within the world of many bio-technological and industrial sectors.

Application of Thermostable Lipases in the Food Industry

Lipases play a large role in the food industry in where they are widely used for manufacturing as well as customization of fats and oils (since they develop an essential element of food) to produce healthier food items. Lipases modify the lipids by causing the alteration within the placement of the important fatty acids in several glycerides [8]. The uniqueness related to lipases ensures they are a perfect option for oleochemical industrial sectors for the manufacturing of several higher value-added products, such as fats in human milk alternatives, the obroma oil counterparts, along with other specific prepared lipids [38]. Since absolutely no method provides such specificities, this amazing trait of thermostable lipases might be focused on industrial and commercial advancements.

Problem/Need

Lipases from *Thermomyces lanuginosus* happen to be identified by producing different transesterification and

esterification reactions. Lipases from *Thermomyces lanuginosus* are recognized to perform the development of glycerides to be able to create healthier kinds of margarine [39]. Lipases seemed to be involved with the growth and development of several favouring agents. Thermostable lipases from several microbes have been employed to result in the oil customization, milk fat manufacturing similar to lipids, production of margarine making use of glycerides, and so on. Modification of fats and oils to produce healthier food products and speciality lipids requires enzymes with high specificity. Chemical catalysis often lacks positional specificity, generates unwanted byproducts, and cannot tailor fatty acid composition effectively.

How Thermostable Lipases Solve It

Thermostable lipases catalyze selective transesterification and esterification reactions, enabling precise alteration of fatty acid positions in glycerides at elevated temperatures, improving stability and reaction rates. This allows the creation of customized lipids, flavor development, and healthier fat substitutes [40].

Examples with Relevant Performance Data

Lipases from *Thermomyces lanuginosus* are recognized for producing modified glycerides used to make healthier margarine and for flavor generation. Beyond classic uses, thermostable lipases assist in synthesizing novel compounds such as mono- and di-acyl esters of glyceryl caffeate with enhanced antioxidant properties [3]. Their effectiveness at elevated temperatures (up to 70°C) supports industrial scalability.

Future Directions

Ongoing industrial innovation aims to improve thermostability and substrate specificity through enzyme engineering [41]. These advances will broaden the application scope in food biotechnology, enabling the production of novel functional lipids, customized fats for nutrition, and improved process efficiency.

Detergent Sector and Thermostable Lipases

The cleaning agent market is among the major marketplaces for enzymes, along with a huge development that takes place to generate the newest enzymes, which include proteases, lipases, and amylases, along with greater as well as significantly better potential. The foremost commercially essential use of Thermostable lipases is within the cleaning agent business, where it's included with the laundry detergent for boosting the second activity [42]. Enzyme revenue in '95 was approximately thirty million dollars, with cleaning agent enzymes creating 30 per cent. The production of laundry detergents is approximately 13 billion tons per year, and the number of lipases is approximately 1,000. One of the primary problems in the laundry sector is the excretion of the adsorbed lipids through the components that are

normally constructed of long-chain, water-insoluble triacylglycerols. These lipids might be eliminated by using cleaners containing lipolytic enzymes, which break them down into complementary fatty acids, diacylglycerols, and monoacylglycerols [43]. The First commercialized lipase used in cleaning agents was lipase TM, which Novo Nordisk introduced in the early nineteen nineties.

Problem/Need

Effective removal of water-insoluble, long-chain triacylglycerol lipid stains under harsh alkaline, high-temperature laundry conditions remains a challenge [44]. Conventional detergents lack sufficient enzymatic components to fully degrade these lipids, impacting cleaning performance and fabric care.

How Thermostable Lipases Solve It

Thermostable lipases hydrolyze complex triacylglycerols into free fatty acids and glycerides that are water-soluble and easily removed. Their stability at alkaline pH and resistance to temperature enable sustained catalytic activity during washing cycles, improving stain removal and preventing fabric greying.

Examples with Relevant Performance Data

Early commercial enzymes such as Lipase TM (originally from a thermophilic fungus, later produced in *Aspergillus oryzae*) and Lumafast™ lipases from *Pseudomonas mendocina* and *P. alcaligenes* demonstrated these properties [37]. Modern developments leverage metagenomic libraries to discover novel, thermostable lipases with enhanced detergent compatibility and stability.

Future Directions

Focused protein engineering and metagenomic screening aim to enhance lipase resistance to surfactants, oxidants, and thermal stress, driving development of detergents with superior cleaning power and reduced environmental impact [45].

Use of Thermostable Lipases in Drug Intermediates and Drugs

Thermostable lipases are also described because of their capability to give rise to the quality of several racemate blends of acids and alcohols [46]. Thermostable lipase, as a result of *Humicola lanuginosa*, has been referred to as among the most effective enzymes for contributing to the differentiation of two enantiomers artificially essential chiral foundation, (Z)-4-triphenyl methoxy-2, epoxy butan-1-ol, using ethylene-vinyl acetate as acyl donor. Lipases from *Thermomyces lanuginosus* demonstrate high enantioselectivity, as shown by their ability to hydrolyze complex bicyclic esters to yield optically pure compounds like (-)-epoxyalcohol and (-)-bromodiol. Additionally, Lipozyme TL IM (a commercial lipase from *T. lanuginosus*) has been effectively used for the regioselective acylation of 5'-O-acyl 5-fluorouracil 1-β-D-ribofuranoside,

enhancing its antitumor potential compared to the parent drug 5-fluorouridine [47]. In pharmaceutical and drug intermediate synthesis, thermostable lipases are gaining prominence for their ability to catalyze chemo-, regio-, and enantioselective reactions, producing high-purity chiral compounds essential for drug development. Their robustness under various solvent conditions makes them ideal for complex organic synthesis routes.

Problem/Need

Stereoselective synthesis of optically pure drug intermediates demands regio-, chemo-, and enantioselective catalysis. Chemical synthesis often produces racemates, lowering drug efficacy and requiring costly purification steps.

How Thermostable Lipases Solve It

Thermostable lipases catalyze selective hydrolysis and acylation reactions even in organic solvents and at elevated temperatures, enabling the preparation of chiral intermediates with high purity and yield.

Examples with Relevant Performance Data

Lipases from *Thermomyces lanuginosus* achieve enantiomeric excess above 96% and purity greater than 99.5% in hydrolyzing bicyclic esters and paclitaxel precursors. Lipozyme TL IM catalyzes regioselective acylation of 5'-O-acyl 5-fluorouracil derivatives, enhancing antitumor potential [48]. *Pseudomonas aeruginosa* lipase demonstrates asymmetric hydrolysis of methoxy-phenyl glycidic acid methyl ester relevant to diltiazem synthesis.

Future Directions

Combining enzyme immobilization with protein engineering is expected to improve reusability, stability, and selectivity [49]. Exploration of thermophilic and extremophilic lipases may yield catalysts suited for broader pharmaceutical applications and more demanding synthesis conditions.

Environmental Applications and Biodegradation

The degradation of atmospheric macromolecules remains a significant challenge; however, this problem can be mitigated to some extent through the application of thermostable lipases. For instance, the side-chains of polyvinyl acetate were successfully hydrolyzed at 60°C in the presence of toluene using various lipases [50]. The efficiency of hydrolysis for long side-chains followed the order: hog pancreatic lipase > Novozym 435 > *Thermomyces lanuginosus* lipase (TLL) > *Candida rugosa* lipase. In contrast, the hydrolysis of shorter side-chains was initiated in the reverse order [40]. The deterioration of several different polymers as an illustration, poly (bisphenol) and polycaprolactone, has been produced at a specific temperature, which ranges between 27 and seventy degrees Celsius, using several lipases, for instance, lipases from hog-pancreas, *Thermomyces lanuginosus*, and Novozyme 436 (N 436) in the existence of

different chemicals [10]. The highest temperature ranges noted for hog pancreas lipases, as well as others, respectively, had been around 50 and 60°C. However, the degradability perspective of lipase action was within the variety of *Thermomyces lanuginosus* > *Candida rugosa* > Novozym-435 > hog pancreas.

Problem/Need

Biodiesel synthesis requires efficient transesterification of lipids; cosmetic industries demand biocompatible, biodegradable polymers synthesized under mild conditions. Enzymes must sustain high temperatures and resist organic solvents [42].

How Thermostable Lipases Solve It

They catalyze lipid transesterification and ring-opening polymerization reactions at elevated temperatures (up to 80°C) and in organic solvents, enhancing process efficiency and product quality [50].

Examples with Relevant Performance Data

Lipase B from *Candida antarctica* and lipases from *Thermomyces lanuginosus* have shown exceptional stability and catalytic performance in biodiesel production and synthesis of polycaprolactone biopolymers used in cosmetics [41]. Their biocompatibility and permeability make them ideal for biomedical applications.

Future Directions

Protein engineering aims to broaden substrate specificity, improve solvent tolerance, and lower production costs [24]. Development of thermostable lipases with tailored catalytic properties will support sustainable biofuel manufacturing and next-generation cosmetic polymers.

CONCLUSION

Because the entire world is focusing on new developments every single day, along with the development in modern technology, the enzymes produced from thermophiles have acquired significant uses and therefore are a source of appeal for a lot of industrial sectors. Thermophilic lipases have grown in popularity during the past several years. Consequently, they are among the essential biocatalysts due to their remarkable capability to accomplish unique interfacial reactions as well as their capacity to catalyze changes. Due to their proficiency in experiencing unpleasant conditions, they've attained immense uses in the areas of bio-technology, pharmaceuticals, and microbiology. Checking up on all the innovative and exclusive characteristics of thermophiles, it's now essential to research completely new bacterial resources for all these enzymes and carefully look for their uses in different areas. Aside from this, methods such as molecular dynamics, as well as developing methods for free energy computation, allow us to obtain an understanding of the elements responsible for thermal resistance.

Authors Contribution

Conceptualization: TB, TH

Methodology: A, SAB, TH, W, IR

Formal analysis: RB

Writing review and editing: TB, TH

All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The authors received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Hamdan SH, Maingwa J, Ali MS, Normi YM, Sabri S, Leow TC. Thermostable Lipases and Their Dynamics of Improved Enzymatic Properties. *Applied Microbiology and Biotechnology*. 2021 Oct;105(19): 7069-94. doi: 10.1007/s00253-021-11520-7.
- [2] Chandra P, Enespa, Singh R, Arora PK. Microbial Lipases and Their Industrial Applications: A Comprehensive Review. *Microbial Cell Factories*. 2020 Aug; 19(1): 169. doi: 10.1186/s12934-020-01428-8.
- [3] Mehta A, Guleria S, Sharma R, Gupta R. The Lipases and Their Applications with Emphasis on Food Industry. In *Microbial Biotechnology in Food and Health*. 2021 Jan: 143-164. doi: 10.1016/B978-0-12-819813-1.00006-2.
- [4] Scully SM and Orlýgsson J. Thermostable Enzymes and Their Applications. In *Thermophilic Anaerobes: Phylogeny, Physiology and Biotechnological Applications*. Cham: Springer International Publishing. 2023 Nov: 155-186. doi: 10.1007/978-3-031-41720-7_6.
- [5] Hussian CH and Leong WY. Thermostable Enzyme Research Advances: A Bibliometric Analysis. *Journal of Genetic Engineering and Biotechnology*. 2023 Dec; 21(1): 37. doi: 10.1186/s43141-023-00494-w.
- [6] Jaiswal N and Jaiswal P. Thermostable α -amylases and Laccases: Paving the Way for Sustainable Industrial Applications. *Processes*. 2024 Jun; 12(7): 1341. doi: 10.3390/pr12071341.
- [7] Ehtiati S and Khatami SH. Lipase: Recombinant Production Methods, Origins, and Industrial Uses. *Biotechnology and Applied Biochemistry*. 2025 May. doi: 10.1002/bab.2781.
- [8] Zhu C, Chen Y, Isupov MN, Littlechild JA, Sun L, Liu X et al. Structural Insights into A Novel Esterase from the East Pacific Rise and Its Improved Thermostability by A Semirational Design. *Journal of Agricultural and Food Chemistry*. 2021 Jan; 69(3):

- 1079-90. doi: 10.1021/acs.jafc.0c06338.
- [9] Abdelaziz AA, Abo-Kamar AM, Elkotb ES, Al-Madboly LA. Microbial Lipases: Advances in Production, Purification, Biochemical Characterization, and Multifaceted Applications in Industry and Medicine. *Microbial Cell Factories*. 2025Feb;24(1):40. doi: 10.1186/s12934-025-02664-6.
- [10] Verma S, Meghwanshi GK, Kumar R. Current Perspectives for Microbial Lipases from Extremophiles and Metagenomics. *Biochimie*. 2021 Mar; 182: 23-36. doi: 10.1016/j.biochi.2020.12.027.
- [11] Sharma D, Bhardwaj KK, Gupta R. Immobilization and Applications of Esterases. *Biocatalysis and Bio-transformation*. 2022May; 40(3):153-68. doi:10.1080/10242422.2021.2013825.
- [12] Lupo A, Haenni M, Madec JY. Antimicrobial Resistance in *Acinetobacter* Spp. and *Pseudomonas* Spp. *Microbiology Spectrum*. 2018Jun;6(3):10-128. doi: 10.1128/microbiolspec.ARBA-0007-2017.
- [13] Seth A. 21 Thermophilic Microbial. *Handbook of Aquatic Microbiology*. 2024Sep;293. doi:10.1201/9781003408543-21.
- [14] Bashir T, Majeed H, Iftikhar T. Isolation and Screening of Thermophilic Bacteria and Its Subsequent Evaluation for Lipases Production. *Pakistan Journal of Botany*. 2024 Apr; 56(2): 759-64. doi: 10.30848/PJ B2024-2(3).
- [15] Masomian M, Rahman RN, Salleh AB, Basri M. A Unique Thermostable and Organic Solvent Tolerant Lipase from Newly Isolated *Aneurinibacillus Thermoaerophilus* Strain HZ: Physical Factor Studies. *World Journal of Microbiology and Biotechnology*. 2010Sep;26(9):1693-701. doi:10.1007/s11274-010-0347-1.
- [16] Peeters C, Cooper VS, Hatcher PJ, Verheyde B, Carlier A, Vandamme P. Comparative Genomics of *Burkholderia Multivorans*, A Ubiquitous Pathogen with A Highly Conserved Genomic Structure. *PLOS One*. 2017Apr;12(4):e0176191. doi:10.1371/journal.pone.0176191.
- [17] Carvalho-Gonçalves LC, Górlach-Lira K. Lipases and Biosurfactants Production by the Newly Isolated *Burkholderia* Sp. *Brazilian Journal of Biological Sciences*. 2018Apr;5(9):57-68. doi:10.21472/bjbs.050906.
- [18] Singh B, Bulusu G, Mitra A. Understanding the Thermostability and Activity of *Bacillus Subtilis* Lipase Mutants: Insights from Molecular Dynamics Simulations. *The Journal of Physical Chemistry B*. 2015 Jan; 119(2): 392-409. doi: 10.1021/jp5079554.
- [19] Konuray G and Erginkaya Z. Potential Use of *Bacillus* Coagulans in the Food Industry. *Foods*. 2018Jun;7(6): 92. doi: 10.3390/foods7060092.
- [20] Balan A. Production, Cloning and Characterization of Thermostable Lipase from *Geobacillus Thermodenitrificans* IBRL-nra (Doctoral dissertation , Universiti Sains Malaysia). 2015.
- [21] Manco G, Adinolfi E, Pisani FM, Ottolina G, Carrea G, Rossi M. Overexpression and Properties of a New Thermophilic and Thermostable Esterase from *Bacillus Acidocaldarius* with Sequence Similarity to Hormone-Sensitive Lipase Subfamily. *Biochemical Journal*. 1998 May; 332(1): 203-12. doi: 10.1042/bj3320203.
- [22] Nawani N, Dosanjh NS, Kaur J. A Novel Thermostable Lipase from A Thermophilic *Bacillus* Sp.: Characterization and Esterification Studies. *Biotechnology Letters*. 1998Oct;20(10):997-1000. doi: 10.1023/A:1005430215849.
- [23] Verma DK, Al-Sahlany ST, Niamah AK, Thakur M, Shah N, Singh S et al. Recent Trends in Microbial Flavour Compounds: A Review on Chemistry, Synthesis Mechanism and Their Application in Food. *Saudi Journal of Biological Sciences*. 2022 Mar; 29(3): 1565-76. doi: 10.1016/j.sjbs.2021.11.010.
- [24] McPherson A, Larson SB, Kalasky A. The Crystal Structures of *Thermomyces* (*Humicola*) *Lanuginosa* Lipase in Complex with Enzymatic Reactants. *Current Enzyme Inhibition*. 2020Dec;16(3):199-213. doi: 10.2174/1573408016999200511090910.
- [25] Abdel-Fattah YR, Gaballa AA. Identification and Over-Expression of A Thermostable Lipase from *Geobacillus Thermoleovorans* Toshki in *Escherichia coli*. *Microbiological Research*. 2008Jan;163(1): 13-20. doi: 10.1016/j.micres.2006.02.004.
- [26] Rath P, Bradoo S, Saxena RK, Gupta R. A Hyper-Thermostable, Alkaline Lipase from *Pseudomonas* Sp. with the Property of Thermal Activation. *Biotechnology Letters*. 2000Mar;22(6):495-8. doi: 10.1023/A:1005604617440.
- [27] Kulkarni N and Gadre RV. A Novel Alkaline, Thermostable, Protease-Free Lipase from *Pseudomonas* sp. *Biotechnology Letters*. 1999Oct; 21(10): 897-9. doi: 10.1023/A:1005591009596.
- [28] Hasan F, Shah AA, Hameed A. Industrial Applications of Microbial Lipases. *Enzyme and Microbial Technology*. 2006Jun;39(2):235-51. doi:10.1016/j.enzmtec.2005.10.016.
- [29] Ikeda M and Clark DS. Molecular Cloning of Extremely Thermostable Esterase Gene from Hyperthermophilic Archaeon *Pyrococcus Furiosus* in *Escherichia coli*. *Biotechnology and Bioengineering*. 1998Mar;57(5):624-9. doi:10.1002/(SICI)1097-0290(19980305)57:5<624::AID-BIT15>3.0.CO;2-B.

- [30] Sun SY, Xu Y, Wang D. Novel Minor Lipase from *Rhizopus Chinensis* During Solid-State Fermentation : Biochemical Characterization and Its Esterification Potential for Ester Synthesis. *Bioresource Technology*. 2009 May; 100(9): 2607-12. doi: 10.1016/j.biortech.2008.11.006.
- [31] Ando S, Ishida H, Kosugi Y, Ishikawa K. Hyperthermostable Endoglucanase from *Pyrococcus Horikoshii*. *Applied and Environmental Microbiology*. 2002 Jan; 68(1): 430-3. doi: 10.1128/AEM.68.1.430-433.2002.
- [32] Iyer PV and Ananthanarayan L. Enzyme Stability and Stabilization—Aqueous and Non-Aqueous Environment. *Process Biochemistry*. 2008 Oct; 43(10): 1019-32. doi: 10.1016/j.procbio.2008.06.004.
- [33] Kumar R, Sharma A, Kumar A, Singh D. Lipase from *Bacillus Pumilus* RK31: Production, Purification and Some Properties. *World Applied Sciences Journal*. 2012; 16(7): 940-8.
- [34] Royter M, Schmidt M, Elend C, Höbenreich H, Schäfer T, Bornscheuer UT et al. Thermostable Lipases from the Extreme Thermophilic Anaerobic Bacteria *Thermoanaerobacter Thermohydrosulfuricus* SOL1 and *Caldanaerobacter Subterraneus* Subsp. *Tengcongensis*. *Extremophiles*. 2009 Sep; 13(5): 769-83. doi: 10.1007/s00792-009-0265-z.
- [35] Kanwar SS, Ghazi IA, Chimni SS, Joshi GK, Rao GV, Kaushal RK et al. Purification and Properties of a Novel Extra-Cellular Thermotolerant Metallolipase of *Bacillus Coagulans* MTCC-6375 Isolate. *Protein Expression and Purification*. 2006 Apr; 46(2): 421-8. doi: 10.1016/j.pep.2005.10.007.
- [36] Shariff FM, Rahman RN, Basri M, Salleh AB. A Newly Isolated Thermostable Lipase from *Bacillus* sp. *International Journal of Molecular Sciences*. 2011 May; 12(5): 2917-34. doi: 10.3390/ijms12052917.
- [37] Li QQ, Zhu ZR, Liu QG, An YT, Wang YX, Zhang SB et al. Characterization of A Novel Thermostable Alkaline Lipase Derived from A Compost Metagenomic Library and Its Potential Application in the Detergent Industry. *Frontiers in Microbiology*. 2022 Dec; 13: 1088581. doi: 10.3389/fmicb.2022.1088581.
- [38] Vivek K, Sandhia GS, Subramaniyan SJ. Extremophilic Lipases for Industrial Applications: A General Review. *Biotechnology Advances*. 2022 Nov; 60: 108002. doi: 10.1016/j.biotechadv.2022.108002.
- [39] Rajak S, Ali SR, Pal B, Chakraborty SS. A Statistical Insight to Exploration of Medicinal Wastewater as A Source of Thermostable Lipase-Producing Microorganisms. *PLoS One*. 2025 Feb; 20(2): e0319023. doi: 10.1371/journal.pone.0319023.
- [40] Cumming H and Marshall SN. Lipase-Catalyzed Synthesis of Mono-And Di-Acyl Esters of Glyceryl Caffeate in Propylene Carbonate and Their Antioxidant Properties in Tuna Oil. *Journal of Biotechnology*. 2021 Jan; 325: 217-25. doi: 10.1016/j.jbiotec.2020.10.021.
- [41] Reyes-Reyes AL, Valero Barranco F, Sandoval G. Recent Advances in Lipases and Their Applications in the Food and Nutraceutical Industry. *Catalysts*. 2022 Aug; 12(9): 960. doi: 10.3390/catal12090960.
- [42] Leykun S, Johansson E, Vetukuri RR, Ceresino EB, Gessesse A. A Thermostable Organic Solvent-Tolerant Lipase from *Brevibacillus* Sp.: Production and Integrated Downstream Processing Using an Alcohol-Salt-Based Aqueous Two-Phase System. *Frontiers in Microbiology*. 2023 Oct; 14: 1270270. doi: 10.3389/fmicb.2023.1270270.
- [43] Ali S, Khan SA, Hamayun M, Lee IJ. The Recent Advances in the Utility of Microbial Lipases: A Review. *Microorganisms*. 2023 Feb; 11(2): 510. doi: 10.3390/microorganisms11020510.
- [44] Darbandi A, Elahi Z, Dadgar-Zankbar L, Ghasemi F, Kakavandi N, Jafari S et al. Application of Microbial Enzymes in Medicine and Industry: Current Status and Future Perspectives. *Future Microbiology*. 2024 Nov; 19(16): 1419-37. doi: 10.1080/17460913.2024.2398337.
- [45] Zhao J, Ma M, Zeng Z, Wan D, Yan X, Xia J et al. Production, Purification, Properties and Current Perspectives for Modification and Application Of Microbial Lipases. *Preparative Biochemistry and Biotechnology*. 2024 Sep; 54(8): 1001-16. doi: 10.1080/10826068.2024.2323196.
- [46] García-Silvera EE, Martínez-Morales F, Bertrand B, Morales-Guzmán D, Rosas-Galván NS, León-Rodríguez R et al. Production and Application of A Thermostable Lipase from *Serratia Marcescens* in Detergent Formulation and Biodiesel Production. *Biotechnology and Applied Biochemistry*. 2018 Mar; 65(2): 156-72. doi: 10.1002/bab.1565
- [47] Ilesanmi OI, Adekunle AE, Omolaiye JA, Olorode EM, Ogunkanmi AL. Isolation, Optimization and Molecular Characterization of Lipase Producing Bacteria from Contaminated Soil. *Scientific African*. 2020 Jul; 8: e00279. doi: 10.1016/j.sciaf.2020.e00279.
- [48] Rozi MF, Rahman RN, Leow AT, Ali MS. Ancestral Sequence Reconstruction of Ancient Lipase from Family I. 3 Bacterial Lipolytic Enzymes. *Molecular Phylogenetics and Evolution*. 2022 Mar; 168: 107381. doi: 10.1016/j.ympev.2021.107381.
- [49] Salihu A and Alam MZ. Thermostable lipases: An Overview of Production, Purification and Characterization. *Biosciences Biotechnology*

Research Asia.2014;11(3):1095-107.doi:10.13005/bbra/1494.

- [50] Mazhar H, Ullah I, Ali U, Abbas N, Hussain Z, Ali SS et al. Optimization of Low-Cost Solid-State Fermentation Media for the Production of Thermostable Lipases Using Agro-Industrial Residues as Substrate in Culture of *Bacillus Amyloliquefaciens*. *Biocatalysis and Agricultural Biotechnology*.2023 Jan; 47: 102559. doi: 10.1016/j.bcab.2022.102559.