



# FUTURISTIC BIOTECHNOLOGY

<https://fbtjournal.com/index.php/fbt>

ISSN (E): 2959-0981, (P): 2959-0973

Vol 05 Issue 01, (Jan-Mar, 2025)



## Review Article



# Genetics of Cotton Fiber Color: Unveiling the Mechanisms and Exploring Gene-Based Approaches for Color Expression

Muhammad Naveed Anjum<sup>1\*</sup>, Muqaddas Amna Qureshi<sup>1</sup>, Maham Riaz<sup>1</sup>, Rohollah Bostani<sup>2</sup>, Azka Ahmad<sup>1</sup> and Nida Shabbir<sup>3</sup>

<sup>1</sup>National Centre of Excellence in Molecular Biology, Punjab University, Lahore, Pakistan

<sup>2</sup>Faculty of Agriculture Sciences, Punjab University, Lahore, Pakistan

<sup>3</sup>Institute of Molecular Biology and Biotechnology, The University of Lahore, Pakistan

## ARTICLE INFO

### Keywords:

Naturally Colored Cotton, Genetics Mechanism, Pigment Biosynthesis, Color Expression

### How to Cite:

Anjum, M. N., Qureshi, M. A., Riaz, M., Bostani, R., Ahmad, A., & Shabbir, N. (2025). Genetics of Cotton Fiber Color: Unveiling the Mechanisms and Exploring Gene-Based Approaches for Color Expression: Genetics of Cotton Fiber: Exploring Gene-Based Approaches. *Futuristic Biotechnology*, 5(1), 21-26. <https://doi.org/10.54393/fbt.v5i1.166>

### \*Corresponding Author:

Muhammad Naveed Anjum  
National Centre of Excellence in Molecular Biology,  
Punjab University, Lahore, Pakistan  
[muhammadnaveedanjum192@gmail.com](mailto:muhammadnaveedanjum192@gmail.com)

Received date: 21<sup>st</sup> January, 2025

Acceptance date: 15<sup>th</sup> March, 2025

Published date: 31<sup>st</sup> March, 2025

## ABSTRACT

Cotton is a vital natural fiber, and its color significantly impacts its commercial value and marketability. Traditional cotton cultivation focuses on white fibers, but naturally colored cotton (NCC) is gaining attention due to its potential for sustainable and environmentally friendly textile production. This review explores the genetics and molecular mechanisms underlying cotton fiber color, focusing primarily on anthocyanins and gossypol pigments. We discuss the role of key genes involved in pigment biosynthesis and regulatory pathways. Additionally, we analyze the potential of gene-based color expression approaches, including traditional breeding and genetic engineering. We highlight the advantages and limitations of each approach and discuss future research directions for developing NCC varieties with improved fiber quality, yield, and color diversity.

## INTRODUCTION

Cotton is one of the most important natural fibers in the world, and its color is a critical quality trait that influences its commercial value and marketability. Traditionally, cotton has been cultivated for its white fibers, but there is growing interest in developing colored cotton varieties for use in various textile applications. Gene-based color expression offers a promising approach to achieving this goal. Throughout the world, in the 1990s, polyester and other man-made fibers faced strong competition. This competition increased further in the early 2000s [1]. Cotton fibers or cotton lint are mostly used in fabrics today [2]. There is a unique way of fiber formation during cotton plant growth. Fiber is created when smaller cells on the

surface of the cotton seed stretch outward. Inner structure breakdown during fiber maturation, and the cell walls shrink, leaving behind a flat, twisted shape [3]. Cotton is a key material in the textile industry. Different dyes are added during production to color fabrics. However, the use of these dyes is causing pollution, which is harmful to human beings [4]. Naturally colored cotton has its color built into the fibers [5, 6]. Scientists are creating improved versions by breeding white cotton with colorful cotton types to produce different colors in cotton fibers naturally. These plants have better color retention, longer fibers, and stronger material than the original colorful cotton. These days, naturally colored cotton is becoming popular




because of the demand for organic, eco-friendly products. Its biggest advantage of naturally colored cotton is that there is no need to add dyes, which is the most polluting step in fabric production, so it is cleaner and safer for both people and the environment [7, 8]. There are many benefits of naturally colored cotton; it serves as a crucial raw material for eco-friendly textiles [9]. It may cut down on chemical residue, pollution, and processing costs. It may also be less flammable and have a higher UV protection rating than regular white cotton [10, 11]. This colored cotton has been grown for many years, but it hasn't improved as fast as regular white cotton. The biggest reasons are that colored cotton plants produce fewer fibers than white cotton, and their colors look faded. These flaws are making them less practical for large-scale farming [9]. Scientists want to fix this by making the fibers stronger, longer and finer [12]. When the cotton bowl opens, the color in the cotton fibers appears. These natural colors come from pigments which are mixed with the cotton's cellulose. This color trait is passed to the next generation through genes. However, traditional breeding struggles to improve both the cotton's quality (like fiber strength) and its yield at the same time; if one gets better, the other might get worse. These days, scientists are using genetic approaches to create these cotton varieties.

Although naturally colored cotton (NCC) offers a sustainable alternative to dyed fabrics, its cultivation faces significant challenges, including low fiber yield, limited color stability, and weaker fiber quality compared to conventional white cotton. While key genes involved in pigment biosynthesis, such as MYB, F3H, and GhANR, have been identified, the molecular regulation of color expression under varying environmental conditions remains insufficiently understood. Moreover, most studies have focused on individual genes or pigment pathways, highlighting a gap in integrative approaches that combine genetics, breeding, and modern gene-editing strategies to achieve both improved fiber quality and consistent color expression.

### Genetics of Cotton Fiber Color

The color of cotton fibers is determined by the accumulation of pigments, primarily anthocyanins and gossypol. These pigments are synthesized through a complex network of biochemical pathways that are regulated by genes. Several genes have been identified that play important roles in cotton fiber color. For example, the MYB113 gene is a transcription factor that regulates the expression of genes involved in anthocyanin biosynthesis. Targeted Mutations in the MYB113 gene can lead to the production of brown or green cotton fibers [13]. Natural compounds called proanthocyanidins (PAs) and related

substances are responsible for the brown color in cotton fibers. Early research found that these flavonoid compounds are responsible for the brown shades [14]. Brown shades, for example, come from compounds made through the plant's natural processes, similar to how flowers or fruits get their colors [15-17]. Pathways by which colored cotton is produced are shown in Figure 1.



**Figure 1:** Genetic pathways in Cotton Fiber Coloration

Cotton gets natural brown or green shades from pigments in the fibers. There is a chemical pathway for the production of pigments. Proanthocyanins pigments are responsible for the brown color, and anthocyanins pigments are responsible for other colors. Gene directly controls the formation of these pigments. MYB, bHLH, and WD40 are important genes that switch to turn on the production of color-related compounds. Different enzymes such as F3H, F3'H, and F3'5'H add chemical groups to naringenin and create different forms of modified naringenin, which results in the production of pigments like pelargonidin (involved in red color formation), cyanidin (involved in magenta color formation), and delphinidin (involved in blue color formation) [18]. In the anthocyanin pathway, one of the important steps of pigments like pelargonidin, cyanidin, and delphinidin formation is the addition of a hydroxyl group to naringenin to create dihydrokaempferol, which gives colors to flowers, and is added by flavanone 3-hydroxylase enzyme., F3H's role is most important in color formation, while other enzymes like F3'5'H modify pigments later [19]. These enzymes play a role in the formation of anthocyanins, which are involved in red, blue, and purple shades in plants [20]. When the F3H enzyme is less active, it changes the color of cotton petals. F3'H and F3'5'H enzymes are involved in red and blue pigments and use the same starting materials in a color production process. Different colors are produced by the rise and fall of the level of these enzymes. For example, Brown cotton fibers have more active GhF3H genes linked to this color process as compared to white fibers [21]. In contrast to white cotton fiber, there is a greater level of

GhCHS, GhC4H, GhF3'H, and GhF3'5'H enzymes in brown cotton fiber expression during flavonoid synthesis. It is studied the expression of the GhCHS, GhANR, and GhLAR genes affects the concentration of anthocyanidin and fiber color [22]. Finding the key genes, their roles in pigment synthesis, and the specific pigments they influence in cotton fiber coloration (Table 1).

**Table 1:** Summarizing Key Genes Involved in Cotton Fiber Coloration

| Gene             | Role in Pigment Synthesis   | Influenced Pigments  | References |
|------------------|---|--|------------|
| MYB113           | Regulates anthocyanin biosynthesis                                | Anthocyanins   | [23]       |
| F3H              | Catalyzes hydroxylation in the anthocyanin pathway                | Dihydroflavonols, leading to pelargonidin, cyanidin, delphinidin | [24]       |
| F3'H             | Involved in flavonoid B-ring hydroxylation for cyanidin synthesis | Cyanidins (red hues)   | [20]       |
| F3'5'H           | Facilitates hydroxylation for delphinidin synthesis               | Delphinidins (blue hues)   | [25]       |
| GhMYB113         | post-anthesis (DPA) fibers with tissue-specific promoter          | brown mature fibers  | [26]       |
| GhPIF4           | Regulate the flowering time of cotton                             | –  | [27]       |
| GhCHS            | Colored fiber formation   | –  | [22]       |
| GhANR, and GhLAR | affect anthocyanidin content and fiber color                      | –  | [22]       |
| GhF3H            | anthocyanin biosynthesis and fiber color formation                | –  | [15]       |
| GhDFR1           | brown color formation of cotton fiber                             | –  | [15]       |
| GhANS            | strongly associated with brown color                              | –  | [29]       |

### Traditional Breeding

This approach involves crossing cotton varieties with different fiber colors to introduce desirable color traits into new varieties. So far, varieties of colored cotton have been developed mostly by selection and recurrent crossing approaches from the germplasm [30]. Selection, the discrimination among variability, is the oldest method of plant improvement [31]. However, this approach is time-consuming and can be challenging due to the complex inheritance patterns of cotton fiber color. The use of statistical genetics and biometrical techniques such as principal component analysis (PCA) and correlation analysis can help in the selection and evaluation of breeding material, optimizing the breeding process, and identifying parental pairs for hybridization [32, 33]. Complex hybridization methods, such as double, complex, and double crossing, have been studied in cotton selection research and have shown promising results in collecting valuable traits from different genetic origins in a single

genotype [34].

### Genetic Engineering

This approach involves directly introducing genes into cotton plants that control the production of pigments. The science of genetics revolutionized the selection process, taking the guesswork out of it, facilitating it and making it more efficient [35]. Modern plant breeders follow standard methods to create variability, discriminate among the variability, and develop cultivars for release to farmers [36]. This approach can be more efficient than traditional breeding methods, but many ethical concerns and regulatory hurdles need to be addressed (Table 2).

**Table 2:** Traditional Breeding and Genetic Engineering

| Aspect                          | Traditional Breeding  | Genetic Engineering  |
|---------------------------------|---|--|
| Time Required                   | Time-consuming due to multiple breeding cycles                          | Faster, direct introduction of desired genes                             |
| Efficiency                      | Variable, depends on genetic variability and inheritance patterns       | Highly efficient, precise targeting of traits                            |
| Challenges                      | Complex inheritance patterns, lower yield of colored fibers             | Regulatory approvals, public acceptance, and possible unintended effects |
| Potential Outcomes              | New varieties with desired color traits and possible yield improvements | Consistent color expression, potential for enhanced fiber qualities      |
| Ethical and Regulatory Concerns | Generally accepted, no significant ethical concerns                     | Ethical concerns require regulatory approvals                            |

### Limitations, Current status and Future Prospects

This study is limited by the variability in genetic expression and environmental influence on fiber pigmentation, which may affect reproducibility across different cotton lines. Future research should focus on integrating advanced genomics, transcriptomics, and gene-editing techniques like CRISPR/Cas9 to stabilize color expression while enhancing fiber quality and yield. Remarkable progress has been made in understanding the genetic basis of cotton fiber color throughout the world. Many key genes have been identified that play important roles in pigment synthesis in cotton, and transgenic cotton lines with altered fiber colors have been developed by using advanced genetic engineering techniques. Large-scale colored cotton production would be possible if we could overcome the following challenges. Improving the color Expression: Improvement in stability and color expression is needed as the color of transgenic cotton fibers can be variable and sensitive to environmental conditions. Ensuring the safety of colored cotton: Colored cotton fibers must be as safe as white cotton fibers and must not cause any health or environmental risks. Addressing regulatory hurdles and ethical concerns: Genetically edited cotton cultivars must pass regulatory, safety and gain regulatory approval before they can be adopted on a large scale. Many more colors in

cotton can be produced once these issues are resolved, and in this way, consumers will be safe from the negative effects which are caused by artificial dyes. This could also lead to new opportunities for textile manufacturers, and it could reduce the extensive cost on use of using dyes.

## CONCLUSIONS

If naturally colored cotton varieties are produced, it may be beneficial for both the textile industry and the environment. By understanding the genetics of cotton fiber colors, their formation and applying advanced breeding and genetic engineering such as CRISPR Cas9 and RNAi techniques, scientists can overcome the limitations of traditional methods and can produce a wider spectrum of sustainable colors. Naturally colored cotton has the potential to reduce environmental pollution which are associated with dyeing processes, in this way it may be the eco-friendly solution for the fashion industry. Not only can above mentioned benefits be obtained from them they also have other qualities as well, such as improved flammability resistance and UV protection. Once these issues are resolved and new genetic engineering techniques are used, gene-based color expression can transform the cotton industry so in the future, textiles will be eco-friendly.

## Authors Contribution

Conceptualization: MNA

Methodology: MNA, MAQ, MR, RB, AA, NS

Formal analysis: RB, AA

Writing and Drafting: MAQ, MR, NS

Review and Editing: MAQ, MR, NS, RB, AA, MNA, NS

All authors approved the final manuscript and take responsibility for the integrity of the work.

## Conflicts of Interest

All the authors declare no conflict of interest.

## Source of Funding

The authors received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Barros MA, Silva CR, Lima LM, Farias FJ, Ramos GA, Santos RC. A Review On Evolution of Cotton in Brazil: GM, White, and Colored Cultivars. *Journal of Natural Fibers*. 2022 Jan; 19(1): 209-21. doi: 10.1080/15440478.2020.1738306.
- [2] Buainain AM, Alves E, Silveira JM, Navarro Z. The Rural World in 21st Century Brazil. The Formation of a New Agrarian and Agricultural Pattern. Embrapa. 2014.
- [3] Lee JA and Fang DD. Cotton as a World Crop: Origin, History, and Current Status. *Cotton*. 2015 Aug; 57: 1-23. doi: 10.2134/agronmonogr57.2013.0019.
- [4] Weisburger JH. Comments on the History and Importance of Aromatic and Heterocyclic Amines in Public Health. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*. 2002 Sep; 506: 9-20. doi: 10.1016/S0027-5107(02)00147-1.
- [5] Nagarajan S, Purushothaman Y, Selvavinayagam M, Govindharaj P, Musthafa A. Studies on Colored Cotton: Biochemical and Genetic Aspects. In *Cotton*. 2022 Oct. doi: 10.5772/intechopen.104898.
- [6] Yan Q, Wang Y, Li Q, Zhang Z, Ding H, Zhang Y *et al.* Up-regulation of Gh TT 2- 3A in Cotton Fibres During Secondary Wall Thickening Results in Brown Fibers with Improved Quality. *Plant Biotechnology Journal*. 2018 Oct; 16(10): 1735-47. doi: 10.1111/pbi.12910.
- [7] Rathinamoorthy R and Parthiban M. Colored Cotton: Novel Eco-Friendly Textile Material for the Future. *Handbook of Ecomaterials*. 2019; 3: 1499-519. doi: 10.1007/978-3-319-68255-6\_91.
- [8] Günaydin GK, Avinc O, Palamutcu S, Yavas A, Soydan AS. Naturally Colored Organic Cotton and Naturally Colored Cotton Fiber Production. *Organic Cotton: Is it a Sustainable Solution?* 2019: 81-99. doi: 10.1007/978-981-10-8782-0\_4.
- [9] Lv YP, Zhao G, Xie YF, Owusu AG, Wu Y, Gao JS. Transcriptome and Metabolome Profiling Unveil Pigment Formation Variations in Brown Cotton Lines (*Gossypium Hirsutum* L.). *International Journal of Molecular Sciences*. 2023 Mar; 24(6): 5249. doi: 10.3390/ijms24065249,
- [10] Crews PC and Hustvedt G. The Ultraviolet Protection Factor of Naturally-Pigmented Cotton. 2005.
- [11] Kibria G, Repon MR, Hossain MF, Islam T, Jalil MA, Aljabri MD *et al.* UV-Blocking Cotton Fabric Design for Comfortable Summer Wears: Factors, Durability and Nanomaterials. *Cellulose*. 2022 Sep; 29(14): 7555-85. doi: 10.1007/s10570-022-04710-7.
- [12] Naoumkina M, Thyssen GN, Fang DD, Jenkins JN, McCarty JC, Florane CB. Genetic and Transcriptomic Dissection of the Fiber Length Trait from A Cotton (*Gossypium Hirsutum* L.) MAGIC Population. *BMC Genomics*. 2019 Dec; 20: 1-4. doi: 10.1186/s12864-019-5427-5.
- [13] Peng Z, Gao Q, Luo C, Gong W, Tang S, Zhang X *et al.* Flavonoid Biosynthetic and Starch and Sucrose Metabolic Pathways Are Involved in the Pigmentation of Naturally Brown-Colored Cotton Fibers. *Industrial Crops and Products*. 2020 Dec; 158: 113045. doi: 10.1016/j.indcrop.2020.113045.

- [14] Shi S, Tang R, Hao X, Tang S, Chen W, Jiang C et al. Integrative Transcriptomic and Metabolic Analyses Reveal That Flavonoid Biosynthesis Is the Key Pathway Regulating Pigment Deposition in Naturally Brown Cotton Fibers. *Plants*. 2024 Jul; 13(15): 2028. doi: 10.3390/plants13152028.
- [15] Zheng H, Duan B, Yuan B, Chen Z, Yu D, Ke L et al. Flavanone and Flavonoid Hydroxylase Genes Regulate Fiber Color Formation in Naturally Colored Cotton. *The Crop Journal*. 2023 Jun; 11(3): 766-73. doi: 10.1016/j.cj.2022.10.004.
- [16] Mei J, Niu Q, Xu K, Huang Y, Bai S, Zhu J et al. GhmiR858 Inhibits the Accumulation of Proanthocyanidins by Targeting GhTt21 in Cotton (*Gossypium Hirsutum*). *Journal of Agricultural and Food Chemistry*. 2023 Oct; 71(41): 15341-51. doi: 10.1021/acs.jafc.3c03884.
- [17] Chen W, Zhang M, Zhang G, Li P, Ma F. Differential Regulation of Anthocyanin Synthesis in Apple Peel Under Different Sunlight Intensities. *International Journal of Molecular Sciences*. 2019 Dec; 20(23): 6060. doi: 10.3390/ijms20236060.
- [18] Wang R, Ren C, Dong S, Chen C, Xian B, Wu Q et al. Integrated Metabolomics and Transcriptome Analysis of Flavonoid Biosynthesis in Safflower (*Carthamus Tinctorius* L.) with Different Colors. *Frontiers in Plant Science*. 2021 Jul; 12: 712038. doi: 10.3389/fpls.2021.712038.
- [19] Yang H, An W, Gu Y, Peng J, Jiang Y, Li J et al. Integrative Metabolomic and Transcriptomic Analysis Reveals the Mechanism of Specific Color Formation in Phoebe Zhennan Heartwood. *International Journal of Molecular Sciences*. 2022 Nov; 23(21): 13569. doi: 10.3390/ijms232113569.
- [20] Liang CY, Rengasamy KP, Huang LM, Hsu CC, Jeng MF, Chen WH et al. Assessment of Violet-Blue Color Formation in *Phalaenopsis* Orchids. *BioMed Central Plant Biology*. 2020 Dec; 20: 1-6. doi: 10.1186/s12870-020-02402-7.
- [21] Khan AQ, Li Z, Ahmed MM, Wang P, Zhang X, Tu L. Eriodictyol Can Modulate Cellular Auxin Gradients to Efficiently Promote in Vitro Cotton Fibre Development. *BioMed Central Plant Biology*. 2019 Dec; 19: 1-5. doi: 10.1186/s12870-019-2054-x.
- [22] Gao J, Shen LI, Yuan J, Zheng H, Su Q, Yang W et al. Functional Analysis of GhCHS, GhANR and GhLAR in Colored Fiber Formation of *Gossypium Hirsutum* L. *BioMed Central Plant Biology*. 2019 Dec; 19: 1-8. doi: 10.1186/s12870-019-2065-7.
- [23] Zhou L, He Y, Li J, Liu Y, Chen H. CBFs Function in Anthocyanin Biosynthesis by Interacting with MYB113 in Eggplant (*Solanum Melongena* L.). *Plant and Cell Physiology*. 2020 Feb; 61(2): 416-26. doi: 10.1093/pcp/pcz209.
- [24] Dai M, Kang X, Wang Y, Huang S, Guo Y, Wang R et al. Functional Characterization of Flavanone 3-Hydroxylase (F3H) and Its Role in Anthocyanin and Flavonoid Biosynthesis in Mulberry. *Molecules*. 2022 May; 27(10): 3341. doi: 10.3390/molecules27103341.
- [25] Qi Y, Lou Q, Quan Y, Liu Y, Wang Y. Flower-Specific Expression of the *Phalaenopsis* Flavonoid 3', 5'-Hydroxylase Modifies Flower Color Pigmentation in *Petunia* and *Lilium*. *Plant Cell, Tissue and Organ Culture*. 2013 Nov; 115: 263-73. doi: 10.1007/s11240-013-0359-2.
- [26] Wang N, Zhang B, Yao T, Shen C, Wen T, Zhang R et al. Re Enhances Anthocyanin and Proanthocyanidin Accumulation to Produce Red Foliated Cotton and Brown Fiber. *Plant Physiology*. 2022 Jul; 189(3): 1466-81. doi: 10.1093/plphys/kiac118.
- [27] Liu LY, Jia MZ, Wang SN, Han S, Jiang J. Identification and Characterization of Cotton Phytochrome-Interacting Factors in Temperature-Dependent Flowering. *Journal of Experimental Botany*. 2023 Jun; 74(12): 3765-80. doi: 10.1093/jxb/erad119.
- [28] Zu QL, Qu YY, Ni ZY, Zheng K, Chen Q, Chen QJ. The Chalcone Isomerase Family in Cotton: Whole-Genome Bioinformatic and Expression Analyses of the *Gossypium Barbadosense* L. Response to *Fusarium* Wilt Infection. *Genes*. 2019 Dec; 10(12): 1006. doi: 10.3390/genes10121006.
- [29] Malik W, Anjum N, Usman Khan M, Abid MA, Ashraf J, Zhang R et al. Genomics of Naturally Colored Cotton: A Way Forward to Initiate Precision Breeding. In *Cotton Precision Breeding*. Cham: Springer International Publishing. 2021 Jun: 173-195. doi: 10.1007/978-3-030-64504-5\_8.
- [30] Majeed S, Chaudhary MT, Mubarik MS, Rana IA, Shaban M, Tan DK et al. Genetics of Biochemical Attributes Regulating Morpho-Physiology of Upland Cotton Under High Temperature Conditions. *Journal of Cotton Research*. 2024 Jan; 7(1): 3. doi: 10.1186/s42397-023-00164-9.
- [31] Younas A, Riaz N, Rashid M, Fiaz S, Tufail A, Noreen Z et al. Modification in Conventional Methods and Modern Plant Breeding Techniques to Enhance Genetic Gain for Future Food Security. *Crop Biofortification: Biotechnological Approaches for Achieving Nutritional Security Under Changing Climate*. 2025 Feb: 377-94. doi: 10.1002/9781394273270.ch22.
- [32] Çakmak F, Çınar VM, Balci Ş, Ünay A. Yield and Fiber Quality Balance in Upland Cotton (*Gossypium Hirsutum* L.) Breeding. *Bangladesh Journal of*

- Botany. 2023 Jun; 52(2): 283-90. doi: 10.3329/bjb.v52i2.67025.
- [33] Madartov B, Abdiev F, Mavlonova N, Shukurov K. Using the Dominance Coefficient in Assessing Cotton Families. In E3S Web of Conferences. 2023; 381(01014). doi: 10.1051/e3sconf/202338101014.
- [34] Kholmurodova G, Barotova A, Namazov S, Yuldasheva R, Jumashv M. Creation of Selected Items with High Fiber Yield and Length Based On Cotton Composite Hybrids. In E3S Web of Conferences. 2023; 371: 01039. EDP Sciences. doi: 10.1051/e3sconf/202337101039.
- [35] Razzaq A, Saleem F, Kanwal M, Mustafa G, Yousaf S, Imran Arshad HM *et al.* Modern Trends in Plant Genome Editing: An Inclusive Review of the CRISPR/Cas9 Toolbox. International Journal of Molecular Sciences. 2019 Aug; 20(16): 4045. doi: 10.3390/ijms20164045.
- [36] Raza A, Mubarik MS, Sharif R, Habib M, Jabeen W, Zhang C *et al.* Developing Drought-Smart, Ready-to-Grow Future Crops. The Plant Genome. 2023 Mar; 16(1): e20279. doi: 10.1002/tpg2.20279.