



Review Article



Keratinophilic Fungi as Eco-Friendly Agents for Poultry Waste Biodegradation: Mechanisms, Applications, and Sustainable Management Strategies

Tehmina Bashir¹, Tahir Hussain², Alamgir³, Adnan Mehmood⁴, Aijaz Ali Kaleri³ and Hira Ilyas⁵¹Department of Botany, Government College University, Lahore, Pakistan²Department of Management Sciences, Hamdard University, Karachi, Pakistan³Department of Health Sciences, Sohail University, Karachi, Pakistan⁴Department of Microbiology, Gulab Devi Educational Complex, Lahore, Pakistan⁵Department of Microbiology, University of Karachi, Karachi, Pakistan

ARTICLE INFO

Keywords:

Biodegradation, Poultry Waste, Sustainable Management, Keratinophilic Fungi

How to Cite:Bashir, T., Hussain, T., Alamgir, ., Mehmood, A., Kaleri, A. A., & Ilyas, H. (2025). Keratinophilic Fungi as Eco-Friendly Agents for Poultry Waste Biodegradation: Mechanisms, Applications, and Sustainable Management Strategies: Keratinophilic Fungi as Eco-Friendly Agents for Poultry Waste Biodegradation. *Futuristic Biotechnology*, 5(4), 03-09. <https://doi.org/10.54393/fbt.v5i4.186>***Corresponding Author:**

Tehmina Bashir

Department of Botany, Government College University, Lahore, Pakistan

Tehminabashir25@gmail.com

Received Date: 10th July, 2025Revised Date: 29th September, 2025Acceptance Date: 6th October, 2025Published Date: 31st December, 2025

ABSTRACT

The rapid expansion of the poultry industry has resulted in the accumulation of large quantities of keratin-rich waste such as feathers, nails, and skin, which pose significant environmental challenges due to their recalcitrant nature. Conventional disposal methods, including incineration and chemical treatment, are energy-intensive and potentially harmful to ecosystems. Keratinophilic fungi offer an eco-friendly alternative through their ability to produce keratinase enzymes that degrade insoluble keratin into valuable amino acids and peptides. This study investigates the biodegradation potential of keratinophilic fungi isolated from poultry waste, focusing on their keratinase production, activity, and potential applications in sustainable waste management. The findings highlight promising fungal strains capable of efficient keratin degradation, underscoring their potential role in circular bioeconomy strategies and industrial applications such as animal feed production, fertilizer formulation, and leather processing.

INTRODUCTION

According to the United Nations, the global human population is projected to reach 10.1 billion by 2060 [1]. Such rapid growth poses serious challenges for humanity, including the persistent issue of hunger. In 2020, approximately 690 million people, around 9% of the global population, were undernourished and lacked access to sufficient and nutritious food [2]. Achieving food security, improving nutrition, and ending hunger are among the core Sustainable Development Goals (SDGs) [3]. However, factors such as climate variability, pest outbreaks, and,

more recently, the COVID-19 pandemic have hindered the establishment of resilient and efficient food systems. One promising approach to addressing these challenges is the promotion of sustainable agricultural practices [4]. As the human population expands, the demand for agricultural products continues to grow, leading to a corresponding increase in agricultural waste. Among these wastes, keratin-rich materials such as poultry feathers are particularly problematic due to their resistance to natural degradation. Presently, most protein-based agro-



industrial by-products are disposed of through incineration, landfilling, or chemical hydrolysis. While these methods are effective for waste reduction, they conflict with sustainable economy principles, often causing environmental damage and missing opportunities for resource recovery. In contrast, biodegradation, the use of living organisms and their enzymes to decompose waste, offers an eco-friendly alternative. This method not only minimizes environmental impact but also yields valuable hydrolysis products, such as amino acids and peptides, which can be used in fertilizers, animal feed, and other biotechnological applications. Such an approach supports a circular economy model, making waste management both sustainable and economically viable [5]. Although both bacteria and fungi produce keratinases, bacterial strains are often favored in industrial applications due to their faster growth rates and relatively simpler enzyme purification processes. In contrast, fungal keratinases generally exhibit broader substrate specificity and greater resilience under harsh conditions [6]. This review aims to investigate the biodegradation potential of keratinophilic fungi isolated from poultry waste, focusing on their keratinase production, activity, and potential applications in sustainable waste management.

Despite the abundance of keratin-rich poultry waste, its effective disposal remains a major environmental challenge due to the recalcitrant nature of keratin and limitations of conventional methods like incineration or chemical hydrolysis. While microbial keratin degradation offers an eco-friendly alternative, most studies have focused on bacterial keratinases, leaving fungal keratinases particularly from indigenous poultry waste isolates—underexplored. Furthermore, there is limited information on optimizing fungal strains for industrial-scale biodegradation and sustainable valorization of keratin into value-added products.

Keratinophilic Fungi: Biology, Ecology, and Pathogenicity

Keratinophilic fungi are saprotrophic microorganisms capable of degrading highly recalcitrant keratinous structures, including feathers, hair, nails, and skin. Their keratinolytic capacity allows them to utilize keratin as the sole source of carbon and nitrogen. These fungi are classified ecologically into three groups: geophilic, zoophilic, and anthropophilic. Geophilic species inhabit soils rich in keratin debris and are mostly saprophytic. Zoophilic species are associated with animals and can cause infections in keratinized tissues, while anthropophilic species primarily infect humans. Among the notable keratinophilic fungi are members of the genera *Trichophyton*, *Microsporum*, and *Epidermophyton*. Although most species are harmless in their natural habitats, some act as opportunistic pathogens capable of causing dermatophytosis in humans and animals. To

ensure safe application in industrial and environmental processes, several mitigation strategies are recommended. Non-pathogenic or low-risk strains such as *Chrysosporium keratinophilum* should be prioritized. Fungi should be cultivated under controlled conditions, including regulated temperature, pH, and containment, to prevent accidental release. Laboratory and industrial personnel must use appropriate personal protective equipment (PPE) such as gloves, masks, and protective clothing. Residual fungal biomass should be inactivated post-application via autoclaving or chemical sterilization, and adherence to biosafety protocols with routine culture monitoring under BSL-1/2 guidelines is essential. Implementing these measures allows the safe integration of keratinophilic fungi in waste management, enzyme production, and other biotechnological applications without compromising human or animal health [7]. Beyond their biotechnological potential, keratinophilic fungi play a significant ecological role in keratin turnover, contributing to nutrient cycling and the biodegradation of otherwise persistent organic matter [8]. Their ecological versatility has encouraged exploration of diverse environments, including poultry waste, to identify strains capable of efficient keratin degradation.

Keratinolytic Fungi from Different Environments

Chicken-associated microbes, particularly fungi, actinomycetes, and certain bacteria, are effective keratin-degrading organisms due to their production of diverse proteolytic and keratinolytic enzymes [9]. Poultry feathers are rich in nitrogen (15–18% dry weight) and sulfur (2–5% dry weight) and are highly resistant to degradation because of strong disulfide bonds. Only specialized organisms, including dermatophytes, *Chrysosporium* species, and some molds like *Fusarium* spp., can biologically break down keratin [10]. Keratin-degrading dermatophytes include anthropophilic species (*Trichophyton rubrum*), zoophilic species (*Trichophyton verrucosum*), and geophilic species (*Trichophyton terrestre*, *Trichophyton georgie*, *Trichophyton ajelloi*, *Microsporum gypseum*, *M. fulvum*). Geophilic dermatophytes and saprophytic *Chrysosporium* species are widespread in soils enriched with keratinous materials and are also found in bird nests and pellets. Among them, *Chrysosporium keratinophilum* (*Aphanoascus keratinophilus*) is particularly effective, producing keratinolytic and proteolytic enzymes that release nutrients such as ammonia and sulfates [11]. The ecological significance of fungal keratinases lies in their ability to degrade otherwise persistent keratin, contributing to nutrient cycling. Often, this degradation involves a complex interplay of enzymes produced by multiple cooperating microorganisms [12].

Keratin Structure and Degradation Mechanism

There is a huge quantity of keratin waste produced by the poultry processing industry in the form of chicken feathers,

consisting of up to 90 percent keratin protein [13, 14]. Keratin is a very stable, fibrous protein, usually rich in a dense lattice of disulfide-bonded structures and the hydrogen bonds, and hydrophobic interactions altogether, of which it is insoluble and resistant against degradation caused by ordinary proteolytic enzymes [15, 16]. Accumulation of such waste material continuously leads to environmental pollution and requires sustainable strategies of management. Structurally, keratin can be classified into α -keratin, β -keratin, and γ -keratin. α -keratins, found in all vertebrates, possess α -helical secondary structures; β -keratins, predominant in reptiles and birds, form β -sheet structures stabilized by non-covalent interactions, making them especially resistant to hydrolysis [17-19]. The high cysteine content in keratin contributes to its recalcitrance, as cysteine residues form strong disulfide bridges that maintain the rigid protein structure. Certain microorganisms, particularly keratinolytic fungi, have evolved mechanisms to degrade keratin through the secretion of keratinases and associated proteases [11]. Notably, members of the phylum Ascomycota, such as *Onygena corvina*, are recognized for their potent keratin-degrading capabilities [20]. Indigenous fungal isolates from poultry waste, therefore, represent a valuable, yet underexplored, biotechnological resource. Keratin degradation generally proceeds via two key stages: sulfitolysis and proteolysis. In the sulfitolysis phase, disulfide bonds are cleaved either enzymatically (e.g., by sulfide reductases) or chemically (e.g., by reducing agents), converting cysteine residues into more accessible forms and loosening the keratin matrix [21-23]. This destabilization enables the next stage of proteolysis, whereby keratinases cleave the exposed peptide bonds to give rise to soluble peptides and free amino acids [24]. Keratinases may also act synergistically with other accessory enzymes, and crude enzyme preparations may prove to be more efficient in keratin degradation as compared to pure keratinase. The realization of the structural complexity of keratin and the enzyme reactions provoking its degradation is essential in the realization of effective, eco-friendly techniques of converting poultry wastes into useful products like biofertilizers, livestock feed, and bioactive molecules [25]. Subsequently, proteolysis takes place after the staining sulfitolysis whereby the exposed peptide bonds are broken and soluble peptides and free amino acids are released by keratin conductivity. There are frequently greater rates of Keratin degradation by crude enzyme preparations as compared with purified Keratinase alone, indicating that there is an interaction between Keratinase and other accessory enzymes. This two-step enzymatic process is essential for the complete biodegradation of keratin-rich materials [26]. Understanding these mechanisms also informs the

strategies for isolating and screening keratinophilic fungi capable of producing potent keratinases, thereby bridging laboratory studies with practical applications.

Isolation and Screening of Keratinophilic Fungi

Keratinophilic fungi are specialized microorganisms capable of degrading keratin, the highly stable structural protein found in feathers, hair, nails, and other keratinized tissues. Their enzymatic keratinolytic activity enables the breakdown of keratin into amino acids and peptides, making them valuable for biotechnological applications, including poultry waste management [27, 28]. These fungi are commonly isolated from soils, poultry litter, and keratin-rich environments using keratin baiting techniques, where materials such as hair, feathers, or nails serve as selective substrates. Dominant genera include *Chrysosporium*, *Trichophyton*, *Microsporum*, *Aspergillus*, and *Penicillium* [29, 30]. Strains are screened for keratinase activity using keratin agar or quantitative feather degradation assays, and potent isolates are selected for further studies under controlled fermentation conditions. Several studies have demonstrated the potential of keratinophilic fungi in poultry waste biodegradation. For instance, Shestakova et al. [5] highlighted the environmental advantages of microbial keratin degradation over conventional disposal methods like incineration or landfilling, emphasizing its role in sustainable waste management and circular economy practices.

Biodegradation of Waste: Microbial Approaches

Building upon laboratory studies of microbial keratin degradation, these fungi can be further utilized in industrial and agricultural settings to transform poultry feather waste into value-added products such as animal feed, fertilizers, and bioactive compounds. Poultry feathers are a significant by-product of the poultry industry, comprising about 5-7% of chicken weight and accounting for several million tons of waste generated annually [32]. These wastes are composed of approximately 90% keratin, a structural protein characterized by high stability due to disulfide bonds [33]. While feathers possess desirable properties such as warmth retention, sound insulation, flexibility, and low density, their accumulation in landfills poses environmental concerns due to their resistance to natural degradation [34, 35]. Microbial degradation of feathers offers an eco-friendly solution for waste management. Various microorganisms, particularly keratinolytic fungi, can hydrolyze keratin into soluble proteins and amino acids by producing keratinase and protease enzymes [11]. These fungi play a crucial role in converting recalcitrant keratin into value-added products, with potential applications in agriculture, biotechnology, and other industries that utilize keratin-containing raw materials [27]. The Ascomycetes group, including species

such as *Onygena corvina*, is well known for its keratin-degrading abilities [21]. Despite promising results, the practical application of fungal biodegradation faces several hurdles. The efficiency of degradation can be influenced by environmental factors such as temperature,

pH, and moisture content. Additionally, the degradation products may be toxic or accumulate in the environment, posing further ecological risks. The economic viability of large-scale fungal biodegradation processes also remains a significant challenge [36] (Table 1).

Table 1: Common Keratinophilic Fungi Isolated from Poultry Waste and their Keratinolytic Properties

Fungal Genus / Species	Sources	Keratinase Activity	Optimum pH	Optimum Temp (°C)	References
<i>Scopulariopsis brevicaulis</i>	Poultry farm soil, feathers	~3.2 KU/mL; ~79% feather degradation	alkaline (≈ 8)	~35°C*	[37]
<i>Trichophyton mentagrophytes</i>	Poultry soil, feather waste	~2.7 KU/mL; ~72% feather degradation	~7.0–8.0	~30–40°C	[38]
<i>Aspergillus flavus</i> (strain K-03)	Feather meal compost/medium	Moderate to high; purified keratinase active up to 2.39× purification, broad activity	~8.0	~45°C	[39]
<i>Chrysosporium keratinophilum</i>	Poultry litter-enriched soil	Potent keratinase producer under submerged culture†	~7–9	mesophilic (28–37°C)	[40]

Applications of Keratinolytic Biodegradation in Poultry Waste Management

Wastes generated from livestock industries such as poultry, swine, and cattle farming are rich in keratin. Among these, chicken feathers represent the largest source, accounting for nearly 9% of a bird's total body weight, with global production estimated at approximately 11.85 million tonnes annually. Despite this abundance, most feathers are discarded, and only a small proportion is reprocessed into products such as feather meal or fertilizers. Improper disposal can make feather residues reservoirs of pathogenic microorganisms, including *Salmonella* spp., *Staphylococcus* spp., and *Clostridium* spp., as well as carriers of veterinary drugs, antibiotics, and chemical contaminants, all of which contribute to soil and water pollution [41]. Current feather disposal practices, such as incineration, eliminate waste but release greenhouse gases and prevent resource recovery [42]. Other non-biodegradation methods similarly pose environmental drawbacks and fail to align with sustainable waste management goals. In contrast, biodegradation using microbial keratinases presents a green alternative, transforming feather waste into value-added products such as fertilizers, animal feed additives, bioelectricity, and biofuels [43]. Both bacteria and fungi produce Keratinases (EC 3.4.21/24/99); while initially linked mainly to pathogenic dermatophytes, non-pathogenic keratinolytic strains have since been identified [44]. Using poultry feathers as a growth substrate for keratinase-producing microorganisms not only lowers enzyme production costs but also supports circular economy practices and reduces the environmental footprint of poultry farming [45].

Limitations and Future prospects

Although keratinophilic fungi show promising biodegradation potential, challenges remain in scaling up processes, ensuring consistent enzyme activity under variable environmental conditions, and managing potential

pathogenicity of certain strains. Future research should focus on developing genetically safe or non-pathogenic fungal strains, optimizing fermentation and enzymatic conditions, and integrating fungal keratin degradation with other biotechnological approaches to enhance efficiency and expand applications in sustainable animal feed, biofertilizers, and circular bioeconomy practices.

CONCLUSION

Keratinophilic fungi isolated from poultry waste represent a promising and eco-friendly solution for the biodegradation of highly recalcitrant keratinous materials. Through their keratinase activity, these fungi can efficiently convert feathers, nails, and other keratin-rich waste into valuable by-products such as amino acids, peptides, and biofertilizers, supporting both environmental sustainability and circular bioeconomy strategies. Beyond waste valorization, their application can reduce reliance on energy-intensive or chemically harsh disposal methods, mitigating environmental pollution and greenhouse gas emissions. However, the presence of opportunistic pathogenic strains highlights the need for careful strain selection, controlled cultivation, and strict adherence to biosafety protocols to prevent human or animal health risks. Future research should focus on optimizing fermentation and degradation conditions, developing non-pathogenic or genetically safe strains, and integrating keratinophilic fungi into large-scale industrial processes. Moreover, combining fungal biodegradation with other biotechnological approaches such as microbial consortia or enzyme engineering may enhance efficiency and expand the range of feasible applications, including sustainable animal feed, biofertilizers, and novel bioproducts. Ultimately, strategic deployment of keratinophilic fungi offers a scalable, environmentally responsible pathway to transform poultry waste into high-value resources.

Authors' Contribution

Conceptualization: TB, TH

Methodology: TB, A, AAK, HI

Formal analysis: TB, AM

Writing and Drafting: TB, TH, AM

Review and Editing: TB, TH, AM, A, AAK, HI

All authors approved the final manuscript and take responsibility for the integrity of the work.

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The authors received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] United Nations, Department of Economic and Social Affairs, Population Division. World Population Prospects 2019: Probabilistic Population Projections [Internet]. 2019. Available from: <https://population.un.org/wpp/Download/Probabilistic/Population/>
- [2] Food and Agriculture Organization of the United Nations. The State of Food and Agriculture 2019: Moving Forward on Food Loss and Waste Reduction. Rome: FAO; 2019. Available from: <http://www.fao.org/documents/card/en/c/ca9699en>
- [3] Gulseven O, Al Harmoodi F, Al Falasi M, Alshomali I. How the COVID-19 Pandemic will Affect the UN Sustainable Development Goals? SSRN Electronic Journal. 2020 May. doi: 10.2139/ssrn.3592933.
- [4] Paudel D, Neupane RC, Sigdel S, Poudel P, Khanal AR. COVID-19 Pandemic, Climate Change, and Conflicts on Agriculture: A Trio of Challenges to Global Food Security. Sustainability. 2023 May; 15(10): 8280. doi: 10.3390/su15108280.
- [5] Shestakova A, Timorshina S, Osmolovskiy A. Biodegradation of Keratin-Rich Husbandry Waste as a Path to Sustainable Agriculture. Sustainability. 2021 Aug; 13(16): 8691. doi: 10.3390/su13168691.
- [6] JL BJ, Dhanasingh I. An Update on Thermostable Keratinases for Protein Engineering Against Feather Pollutants. Applied Microbiology and Biotechnology. 2025 Dec; 109(1): 1-22. doi: 10.1007/s00253-025-13459-5.
- [7] Pasrija R and Kumari D. Protocols for Studying Pathogenic Fungi. Springer; 2025 May. doi: 10.1007/978-981-96-7047-5.
- [8] Khan AM, Bhadauria S, Yadav R. Keratinophilic Fungi: Isolation, Identification, Pathogenicity, Characterization, and Treatment. In: New and Future Developments in Microbial Biotechnology and Bioengineering. Elsevier. 2020 Jan. p. 29-37. doi: 10.1016/B978-0-12-821006-2.00003-0.
- [9] Tamreihao K, Mukherjee S, Khunjamayum R, Devi LJ, Asem RS, Ningthoujam DS. Feather Degradation by Keratinolytic Bacteria and Biofertilizing Potential for Sustainable Agricultural Production. Journal of Basic Microbiology. 2019 Jan; 59(1): 4-13. doi: 10.1002/jobm.201800434.
- [10] Calin M, Constantinescu-Aruxandei D, Alexandrescu E, Raut I, Doni MB, Arsene ML, et al. Degradation of Keratin Substrates by Keratinolytic Fungi. Electronic Journal of Biotechnology. 2017 Jul; 28: 101-12. doi: 10.1016/j.ejbt.2017.05.007.
- [11] Bohacz J. Biodegradation of Feather Waste Keratin by a Keratinolytic Soil Fungus of the Genus Chrysosporium and Statistical Optimization of Feather Mass Loss. World Journal of Microbiology and Biotechnology. 2017 Dec; 33(1): 13. doi: 10.1007/s11274-016-2177-2.
- [12] Neemisha and Sharma S. Soil Enzymes and Their Role in Nutrient Cycling. In: Structure and Functions of Pedosphere. Singapore: Springer Nature Singapore. 2022 Jun. p. 173-188. doi: 10.1007/978-981-16-8770-9_8.
- [13] Kumawat TK, Sharma A, Sharma V, Chandra S. Keratin Waste: The Biodegradable Polymers. In: Keratin. IntechOpen. 2018 Dec.
- [14] Maurya SD and Singh A. Application and Future Perspectives of Keratin Protein Extracted from Waste Chicken Feather: A Review. Sustainable Chemical Engineering. 2024: 31-45.
- [15] Donato RK and Mija A. Keratin Associations with Synthetic, Biosynthetic and Natural Polymers: An Extensive Review. Polymers. 2019 Dec; 12(1): 32. doi: 10.3390/polym12010032.
- [16] Shavandi A, Silva TH, Bekhit AA, Bekhit AE. Keratin: Dissolution, Extraction, and Biomedical Application. Biomaterials Science. 2017; 5(9): 1699-1735. doi: 10.1039/C7BM00411G.
- [17] Kadir M, Wang X, Zhu B, Liu J, Harland D, Popescu C. The Structure of the "Amorphous" Matrix of Keratins. Journal of Structural Biology. 2017 May; 198(2): 116-23. doi: 10.1016/j.jsb.2017.04.001.
- [18] Strasser B, Mlitz V, Hermann M, Tschachler E, Eckhart L. Convergent Evolution of Cysteine-Rich Proteins in Feathers and Hair. BMC Evolutionary Biology. 2015 May; 15(1): 82. doi: 10.1186/s12862-015-0360-y.
- [19] Passos BM, Rocha VH, Pacheco MS, Perrechil F, Viganó J, de Moraes MA. Extraction, properties, and applications of keratin-based films and blends. In Handbook of Natural Polymers. 2023 Jan; 1: 399-420. Elsevier. doi: 10.1016/B978-0-323-99853-6.00025-5.

- [20] Huang Y, Busk PK, Herbst FA, Lange L. Genome and Secretome Analyses Provide Insights into Keratin Decomposition by Novel Proteases from the Non-Pathogenic Fungus *Onygena Corvina*. *Applied Microbiology and Biotechnology*. 2015 Nov; 99(22): 9635-49. doi: 10.1007/s00253-015-6805-9.
- [21] Peng Z, Zhang J, Du G, Chen J. Keratin Waste Recycling Based on Microbial Degradation: Mechanisms and Prospects. *ACS Sustainable Chemistry and Engineering*. 2019 May; 7(11): 9727-36. doi: 10.1021/acssuschemeng.9b01527.
- [22] Tantamacharik T. Microbial Proteases from Gelatin Production System for the Degradation of Keratin-Containing Materials [dissertation]. Dunedin: University of Otago. 2021
- [23] Das S, Das A, Das N, Nath T, Langthasa M, Pandey P, et al. Harnessing the Potential of Microbial Keratinases for Bioconversion of Keratin Waste. *Environmental Science and Pollution Research*. 2024 Oct; 31(46): 57478-507. doi: 10.1007/s11356-024-34233-6.
- [24] Ardyati T. Screening of Keratinolytic Fungi for Biodegradation Agent of Keratin from Chicken Feather Waste. IOP Conference Series: Earth and Environmental Science. 2019 Dec; 391(1): 012027. doi: 10.1088/1755-1315/391/1/012027.
- [25] Li Q. Progress in Microbial Degradation of Feather Waste. *Frontiers in Microbiology*. 2019 Dec; 10: 2717. doi: 10.3389/fmicb.2019.02717.
- [26] Mercer DK and Stewart CS. Keratin Hydrolysis by Dermatophytes. *Medical Mycology*. 2019 Jan; 57(1): 13-22. doi: 10.1093/mmy/myx160.
- [27] Ojha S. Keratin-Baiting Technique: Method for Isolation of the Keratinophilic Fungi from Soil. *International Journal of Creative Research Thoughts*. 2025 Jan.
- [28] Hamada N, Baba T, Sakuma D. Culturable Keratinophilic Fungi Isolated from House Dust by Hair-Baiting Technique. *International Biodeterioration and Biodegradation*. 2024 Jan; 186: 105702. doi: 10.1016/j.ibiod.2023.105702.
- [29] Sharma R. Isolation and Molecular Characterization of Keratinophilic Fungi. *Research Journal of Science and Technology*. 2024 Jul; 16(3): 181-8. doi: 10.52711/2349-2988.2024.00027.
- [30] Spychala K, Klosinska K, Salwinska W, Ogorek R. Diversity of Soil-Borne Fungi Isolated from Places Frequently Visited by People in the City of Wroclaw (Poland). *Applied Sciences*. 2024 Mar; 14(7): 2782. doi: 10.3390/app14072782.
- [31] Amuthavalli T and Ravi C. Chicken Feather Degradation by Keratinolytic *Aspergillus* Species Isolated from Feather Waste Dumping Soil. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences*. 2022; 24(3): 459-67. doi: 10.53550/AJMBES.2022.v24i03.003.
- [32] Tesfaye T, Sithole B, Ramjugernath D, Chunilall V. Valorization of Chicken Feathers: Characterization of Chemical Properties. *Waste Management*. 2017 Oct; 68: 626-35. doi: 10.1016/j.wasman.2017.06.050.
- [33] Mattiello S, Guzzini A, Del Giudice A, Santulli C, Antonini M, Lupidi G, et al. Physico-Chemical Characterization of Keratin from Wool and Chicken Feathers Extracted Using Refined Chemical Methods. *Polymers*. 2022 Dec; 15(1): 181. doi: 10.3390/polym15010181.
- [34] Alsamaraie AR, Samin PM, Mazali II, Jadallah AA, Sultan KF. Insulation Materials Based on Recycled Feather Waste: A Review. *Tikrit Journal of Engineering Sciences*. 2023 Mar; 30(1): 104-11. doi: 10.25130/tjes.30.1.10.
- [35] Suresh S, Arisutha S, Chaturvedi K, Verma S. Waste Feathers as a Resource, Its Availability, and the Recent Technology in the Production of New Thermal Insulation Materials. In: *Development of Sustainable Thermal Insulators from Waste Materials: A Circular Economy Approach*. Singapore: Springer Nature Singapore; 2024 Sep; 139-47. doi: 10.1007/978-981-97-5444-1_9.
- [36] Vaksmaa A, Guerrero-Cruz S, Ghosh P, Zeghal E, Hernando-Morales V, Niemann H. Role of Fungi in Bioremediation of Emerging Pollutants. *Frontiers in Marine Science*. 2023 Mar; 10: 1070905. doi: 10.3389/fmars.2023.1070905.
- [37] Anbu P, Gopinath SC, Hilda A, Mathivanan N, Annadurai G. Secretion of Keratinolytic Enzymes and Keratinolysis by *Scopulariopsis brevicaulis* and *Trichophyton mentagrophytes*: Regression Analysis. *Canadian Journal of Microbiology*. 2006 Nov; 52(11): 1060-9. doi: 10.1139/w06-067.
- [38] Anbu P, Hilda A, Sur HW, Hur BK, Jayanthi S. Extracellular Keratinase from *Trichophyton* sp. HA-2 Isolated from Feather Dumping Soil. *International Biodeterioration and Biodegradation*. 2008 Oct; 62(3): 287-92. doi: 10.1016/j.ibiod.2007.07.017.
- [39] Kim JD. Purification and Characterization of a Keratinase from a Feather-Degrading Fungus, *Aspergillus flavus* Strain K-03. *Microbiology*. 2007 Dec; 35(4): 219-25. doi: 10.4489/MYCO.2007.35.4.219.
- [40] Al-Bader SM. Proteolytic Activity and Health Impact of Fungi Isolated from Chickens' Feathers in House Breeding Cages. *Iraqi Journal of Agricultural Sciences*. 2023; 54(2): 579-88.
- [41] Cornejo J, Pokrant E, Carvallo C, Maddaleno A, San Martin B. Depletion of Tylosin Residues in Feathers,

- Muscle, and Liver from Broiler Chickens After Completion of Antimicrobial Therapy. *Food Additives and Contaminants: Part A*. 2018 Mar; 35(3): 448-57. doi: 10.1080/19440049.2017.1401740.
- [42] Kodak S, Gharge T, Chavan V. Microbial Degradation of Poultry Feather Wastes under the Influence of Temperature and pH: A Review. *International Journal of Environmental Sciences and Natural Resources*. 2019; 21(3): 87-92. doi: 10.19080/IJESNR.2019.21.556063.
- [43] Peng Z, Mao X, Zhang J, Du G, Chen J. Biotransformation of Keratin Waste to Amino Acids and Active Peptides Based on Cell-Free Catalysis. *Biotechnology for Biofuels*. 2020 Apr; 13(1): 61. doi: 10.1186/s13068-020-01700-4.
- [44] Aktayeva S and Khassenov B. High Keratinase and Other Types of Hydrolase Activity of the New Strain of *Bacillus paralicheniformis*. *Public Library of Science One*. 2024 Oct; 19(10): e0312679. doi: 10.1371/journal.pone.0312679.
- [45] Al Subhi AH. Microbial Keratinase Production Using Chicken Feather Waste. Muscat (OM): Sultan Qaboos University. 2022.